

## Article

# Factores de Rendimiento en Escalada Deportiva y Escalada En Bloque: Revisión Sistemática

## Performance Factors in Sport Climbing and Bouldering: Systematic Review

Isaac López Laval y Sebastian Sitko

Universidad de Zaragoza. Grupos de Investigación del Movimiento Humano - Human Research

### RESUMEN

**Objetivos:** Hasta la fecha no se ha realizado una revisión sistemática sobre esta temática por lo que este documento pretende analizar toda la información disponible sobre los factores de rendimiento en la escalada deportiva y escalada en bloque y proporcionar un documento basado en la ciencia a los atletas, entrenadores e investigadores.

**Métodos:** Se realizó una búsqueda en cuatro bases de datos con las palabras "Sport Climbing" y "Athletic Performance" así como "Bouldering" y "Athletic Performance". Los criterios de inclusión y exclusión incluían estudios realizados con hombres y mujeres adultos y adolescentes sanos que incluyeran un estudio de los distintos factores de rendimiento en la escalada deportiva y escalada en bloque.

**Resultados:** 33 estudios cumplieron los criterios. La fuerza específica de los flexores de antebrazo y dedos y la del tronco superior, la capacidad reoxidativa de la musculatura local y un óptimo ratio fuerza-peso son cruciales para el rendimiento. Los factores psicológicos como el control del estrés y la previsualización han sido menos estudiados. La forma cardiorrespiratoria fue similar otros deportes intermitentes y los factores antropométricos no resultaron tan importantes.

**Conclusiones:** Se obtuvieron resultados uniformes en cuanto a la importancia de factores de fuerza, resistencia, cardiorrespiratorios, psicológicos y antropométricos como claves en la mejora del rendimiento.

**Palabras Clave:** Escalada Deportiva, Escalada en Bloque, Rendimiento atlético

### ABSTRACT

**Objectives:** This systematic review aimed to analyze all the available information on the main performance factors in sport climbing and bouldering and to provide a science-based document for coaches, athletes and researchers.

**Methods:** A search within four different databases was conducted with the words "Sport Climbing" and "Athletic Performance" and "Bouldering" and "Athletic Performance". The results of the search were assessed with the inclusion and exclusion criteria, which included all types of published studies performed on healthy adolescent and adult men and women that studied the different performance factors in climbing and bouldering.

**Results:** 33 studies met the inclusion and exclusion criteria. Specific forearm and finger flexor strength, re-oxydation capacity of the local muscles, general upper body strength and an optimal power to weight ratio seem crucial for performance. Psychological factors such as stress control and previsualization may also play an important role. Cardiorespiratory fitness found in climbers is similar to what is usually found in other intermittent sports. Finally, anthropometric factors are not as important as usually reported in general literature.

**Conclusions:** Although several confounding factors were present, there was general agreement upon the importance of several strength, endurance, anthropometric, cardiorespiratory and psychological factors as keys in performance enhancement in sport climbing and bouldering.

**Keywords:** Sport Climbing, Bouldering, Athletic Performance

## INTRODUCTION

---

Climbing is an intermittent sport that has earned significant popularity in the last decades. Its inclusion in the 2020 Tokyo Olympic Games has had a significant impact in the specialization and training methods of the professional athletes that participate in this sporting activity (Lutter, El-Sheikh, Schöffl, & Schöffl, 2017).

The differences between the sports of bouldering and sport climbing are mainly related to the duration of efforts, which can vary between the 30 seconds normally taken to climb a boulder problem (White & Olsen, 2010) and efforts of up to half an hour in sport climbing or other disciplines (Egocheaga et al., 2001).

A previous narrative review (Phillip B Watts, 2004) performed in 2004 established that elite rock climbers presented low stature and body weight, low percentages of body fat, moderate aerobic endurance and high levels of upper body strength, endurance and power to weight ratio. In addition to these findings, absolute finger strength measured with dynamometry in elite climbers was not necessarily higher than in the general population, although weight adjusted measures did show significant differences (Giles, Rhodes, & Taunton, 2006). This characteristic showed the importance of weight in this gravity-dependent sport.

During the last years several improvements have been made in the field of training for climbing. The appearance of climbing treadmills, better climbing gyms and apps for training monitoring has professionalized elite training. This, together with the increasing proximity of the Olympic Games has produced an increase of interest in the key performance factors of these sport modalities and has been linked to an increased scientific production in the last years. No previous systematic review on this topic has been performed and this study attempts to compile the most important data around performance in climbing in order to offer coaches and athletes a summary of the most important findings around their sport discipline.

## METHODS

---

### Search strategy

The project followed the systematic review methodology proposed in the “Preferred Reporting Items for Systematic reviews and Meta-Analyses” (PRISMA) statement (Moher, Liberati, Tetzlaff, Altman, & PRISMA Group, 2009). The identification of studies was performed by searching within PubMed, Sportdiscus, Scopus and Medline. The search was conducted up to and including December 22nd 2018. Two different searches were performed in each database to ensure that all published studies regarding the topic were included in the present systematic review. The first search was performed using the thesaurus provided by each database: Sport Climbing AND Athletic Performance, while the second search was performed with the following combination of terms: Bouldering AND Athletic Performance. Two reviewers independently examined each database to obtain the potential publications. Full texts of the relevant articles were obtained and assessed against the inclusion and exclusion criteria described below. Inter-reviewer disagreements were resolved by consensus.

### Inclusion criteria

The following inclusion criteria were used: types of studies: cross-sectional, longitudinal, randomized and nonrandomized controlled trials studying the climbing performance factors; types of participants: male and female adults and adolescents; types of outcome measured: main climbing-related performance factors.

### Exclusion criteria


The following exclusion criteria were used: studies in languages other than English or Spanish; unpublished data; studies with animals and dissertations or abstracts from society proceedings or congresses and other similar unpublished data.

### Quality assessment

Studies were assessed with the tool Critical appraisal Skills Programme (CASP) in order to control risk of bias of each one of the studies. All of the included studies were accepted by Research Ethical Committees and complied with the ethical rules set in Declaration of Helsinki.

### Data extraction

All the articles were assessed first according to the title, second according to the abstract and lastly a full review of the article was performed. Figure 1 represents the PRISMA flow chart of the process of study selection. Article selection was performed using a continuous string quadrant in Microsoft Excel 2007.



## RESULTS

A review of the studies included in the qualitative analysis is provided. Subcategories were established in order to differentiate between types of performance factors. Basic information about the studies included in the review can be found in Table 1.

Table 1: Main information about the studies included in the qualitative analysis

Study	Participants	Intervention	Outcome
Cardiorespiratory fitness			
Aras et al, 2016	19 adults	Effects of climbing on physical condition values	Climbing improved muscle strength and endurance, VO <sub>2</sub> max and reduced both fat mass and fat percentage.
Balas et al, 2014	26 male climbers	Measurement of physiological variables during a treadmill climbing-specific test.	Increase in heart rate and oxygen consumption were related to the inclination of the wall.
Bertuzzi et al, 2007	6 elite climbers and 7 recreational climbers	Comparison between energetic systems used by both groups while climbing.	Both groups used the same energetic systems. Elite climbers had better climbing economy and thus performed better on stronger wall inclinations.
Bertuzzi et al, 2012	13 climbers, 6 elite and 7 recreational	Comparison of physiological values while climbing.	Elite group presented lower reductions in grip strength and lower oxygen cost per movement.
Billat et al, 1995	4 high level climbers	Measurement of VO <sub>2</sub> consumption, heart rate and blood lactate while climbing two high difficulty routes.	Fractions of VO <sub>2</sub> max, maximal heart rate and blood lactate used during the ascents varied between the steep and the technical routes.
Limonta et al, 2018	7 advanced and 6 elite climbers	Comparison of oxygen demand on a cycling ergometer and a climbing treadmill	VO <sub>2</sub> consumption estimated during cycling incremental tests cannot be extrapolated to VO <sub>2</sub> consumptions during climbing.
Mermier et al, 1997	14 experienced climbers, 9 men and 5 women	Measurement of physiological variables while climbing three routes with different inclinations.	Nonlinear relationship between increase in VO <sub>2</sub> consumption and heart rate during climbing.

Rodio et al, 2008	13 climbers, 8 men and 5 women	Measurement of lactate, energy expenditure and VO <sub>2</sub> consumption while climbing.	Energy expenditure and VO <sub>2</sub> consumption during climbing are similar to values obtained during easy to moderate aerobic activities.
Watts et al, 2000	15 expert rock climbers	Comparison between active and passive recovery between intervals of climbing.	VO <sub>2</sub> consumption reaches a plateau, blood lactate keeps increasing during the effort. VO <sub>2</sub> consumption doesn't reflect total energy demand during climbing.
Study	Participants	Intervention	Outcome
Strength, endurance and anthropometric factors			
Balas et al, 2016	22 climbers	Comparison of different recovery strategies between intervals of climbing.	Hand shaking improved muscle re-oxygenation and optimized recovery between intervals.
Cutts et al, 1993	25, 13 of them climbers.	Comparison between finger strengths	Only pinch grip stronger in climbers. Not correlated to climbing performance.
Deyhle et al, 2015	11 male climbers	Impact of pre fatigue of different muscle groups on performance.	Digit and elbow flexors were the most determinant muscle groups in climbing performance.
Donath et al, 2013	28 climbers, 14 recreational and 14 advanced.	Determination of differences in load application according to the climbing level.	Better climbers demonstrated more symmetrical load application.
Fryer et al, 2015	38 climbers	Comparison between blood flow and muscle oxygenation during climbing specific tasks.	Muscle re-oxygenation didn't depend on blood flow.
Fryer et al, 2016	46 climbers, 36 males and 10 females.	Determination of an oxidative index of the finger flexors as a predictor of performance.	Oxidative capacity in the finger flexors predicted redpoint performance.

Grant et al, 1996	10 elite climbers, 10 recreational climbers and 10 matched controls.	Comparison of key performance indicators	Elite climbers had better finger and shoulder girdle strength and better hip flexibility.
Grant et al, 2001	30 women, 10 elite climbers, 10 recreational and 10 physically active	Comparison of performance factors	Greater finger strength in elite than in recreational and physically active women.
Grant et al, 2003	27, 9 climbers, 9 rowers and 9 leg trained athletes.	Comparison between finger specific strength.	No significant differences observed between the study groups
Laffaye et al, 2016	41 climbers, 15 beginners, 16 advanced and 10 elite.	Measurement of variables that could explain climbing performance	Almost half of the variance is explained by trainable factors while only a small part is explained by genetic factors.
MacLeod et al, 2007	20, 11 intermediate climbers.	Comparison of specific finger endurance.	Muscle re-oxygenation was higher in climbers and was positively correlated to climbing performance.
Mermier et al, 2000	44 climbers, 24 men and 20 women	Measurement of variance that could explain climbing level.	Almost two thirds of the variance in climbing performance could be explained by trainable factors. Anthropometric factors have only a small role in climbing performance.
Ozimek et al, 2017	20, 14 advanced and 6 elite climbers.	Comparison of strength, endurance and anthropometric factors	Elite climbers present better finger strength and arm endurance than advanced climbers.
Philippe et al, 2012	24, 12 elite male and female climbers, 12 physically active matched controls	Comparisons in flexor strengths, muscle endurance and oxygenation	Superior endurance and muscle oxygenation in climbers than in controls.
Schreiber et al, 2015	31 experienced male rock climbers	Evaluation of connective tissue in hands	Increased thickness of tendon flexor pulleys and articular capsule of the distal interphalangeal joint.

Schoffl et al, 2006	28 male climbers	Performance measures taken while climbing on a treadmill.	Moderate lactate accumulation observed and the reduction in heart rate was considered useful as a recovery indicator.
Vigoureux et al, 2015	25, 12 climbers, 9 men and 3 women and 13 non-climbers.	Comparisons in capacities of finger flexors.	Both male and female climbers presented stronger finger flexors than non-climbers.
Wall et al, 2004	18 women, 6 moderate level, 6 advanced and 6 elite climbers.	Performance on 2 different climbing tests and strength evaluations.	Elite climbers presented better hand strength and one arm lock-off strength than advanced and moderate climbers.
Watts et al, 1996	11 expert climbers	Measurement of handgrip strength, endurance and blood lactate evolution after several intervals of maximal intensity climbing.	Both handgrip strength and endurance are reduced after several intervals of climbing. Handgrip strength recovers faster than endurance and blood lactate is still elevated after 20 minutes of complete rest.
Study	Participants	Intervention	Outcome measured
Psychological factors			
Asci et al, 2007	64 climbers	Comparison of sensation seeking, self-perception and motivation in different groups of climbers.	Climbers presented high sensation seeking, internal motivation and positive self-perception regardless of their performance level.
Fryer et al, 2013	21 climbers, 18 men and 3 women	Comparison of stress during top-rope or lead ascents.	High level climbers were not affected by stress during lead climbing.
Hardy et al, 2007	10 experienced climbers	Evaluation of anxiety levels in different groups of climbers.	Higher anxiety levels were positively correlated to performance and higher level of climbing.
Nieuwenhuys et al, 2008	12 non-climbers, 7 men and 5 women.	Evaluation of the effects of anxiety on visual information processing during climbing.	Anxiety produced worse information processing during climbing in non-climbers.
Sanchez et al, 2012	29 male climbers	Evaluate of the effects of route previsualization on climbing performance of different climbing levels.	Only expert climbers did benefit from route previsualization.

## Climbing and cardiorespiratory fitness

One study (Billat, Dalleja, Charlaix, Rizzardo, & Janel, 1995) analyzed the metabolic requirements during the completion of two competitive sport climbing routes and reported that submaximal levels of VO<sub>2</sub>max (37.7 to 45.6 ml/kg/min) were reached together with moderate blood lactate concentrations (4.3 mmol.l<sup>-1</sup> to 5.7 mmol.l<sup>-1</sup>) and high but no maximal heart rate were obtained (77 to 85.5% of maximal heart rate). When the same values were analyzed in highly trained boulderers (La Torre, Crespi, Serpiello, & Merati, 2009) slightly higher values were obtained (6.6 mmol.l<sup>-1</sup>) VO<sub>2</sub> consumption and heart rate have been found to increase with wall inclination until reaching submaximal (VO<sub>2</sub>) and maximal (heart rate) values (Balas et al., 2014). Similar results were obtained (Mermier, Robergs, McMinn, & Heyward, 1997) in another study that also reported a non-linearity between heart rate and VO<sub>2</sub> consumption during rock climbing activities.

These values have also been reported in non-competitive climbers (Rodio, Fattorini, Rosponi, Quattrini, & Marchetti, 2008), with VO<sub>2</sub>max values of around 40 ml/kg/min and oxygen consumption during climbing efforts that didn't surpass 28.3 ml/kg/min. Another study (P B Watts, Daggett, Gallagher, & Wilkins, 2000) reported a VO<sub>2</sub> plateau despite a continuous increase in blood lactate, which would explain a higher energy demand than the one indicated during VO<sub>2</sub> recording during climbing. VO<sub>2</sub> and blood lactate values didn't return to normal even after 20 minutes of complete rest. These results coincide with the findings reported by another study (R. C. de M. Bertuzzi, Franchini, Kokubun, & Kiss, 2007) which reported that the main energy contribution during rock climbing comes from the aerobic and anaerobic alactic systems. Other researchers (Limonta et al., 2018) have suggested that care should be taken while estimating energetic demands based on VO<sub>2</sub> consumption while climbing.

Oxygen cost per movement has also been reported as a reliable tool to measure performance in climbers (R. Bertuzzi et al., 2012). The aerobic component of the sport was proven in an intervention study with sedentary subjects who completed a climbing training intervention. Climbing produced improvements in body composition and increases in VO<sub>2</sub>max, muscle strength and endurance (Aras & Akalan, 2016), findings similar to those obtained with moderate endurance exercise such as running or cycling with sedentary subjects.

### ***Strength, endurance and anthropometric factors in climbing***

A study (Grant, Hynes, Whittaker, & Aitchison, 1996) compared the differences in endurance, strength, flexibility and anthropometry in elite rock climbers, recreational climbers and non-climbers. Elite rock climbers performed a significantly higher number of pull ups and lasted longer during a bent-arm hang. Also, better climbing ability was correlated to increased hip flexibility, shoulder girdle strength and endurance as well as greater finger strength. Increased finger strength in elite climbers was also reported even when this group was compared to advanced climbers (Ozimek et al., 2017). Finger strength in elite climbers is also a result of years of adaptation to climbing stimulus which results in increased thickness of flexor tendon pulleys and capsules of the distal inter-phalangeal joints (Schreiber, Allenspach, Seifert, & Schweizer, 2015).

Similar studies have been performed with female climbers, (Grant et al., 2001) showing again significantly greater finger strength in the elite group compared to the amateur and non-climbing groups. Another study found similar results (Wall, Starek, Fleck, & Byrnes, 2004), where climbing-specific hand strength and one-arm lock off strength adjusted to body weight were higher in elite female climbers than in amateur or non-climbing matched groups and also predicted climbing performance.

One study (Mermier, Janot, Parker, & Swan, 2000) performed a complete anthropometric, physiological and demographic analysis of 44 climbers which were classified by their performance levels. The study concluded that a 58.9% of the total variance in climbing performance could be explained by trainable characteristics, especially strength and endurance. Researchers refuted the belief that specific anthropometric characteristics were needed in order to perform in the sport. These results were verified in another more recent piece of research which found that anthropometric factors explained only a 4% of climbing performance while trainable variables explained a 46% (Laffaye, Levernier, & Collin, 2016).

An old study (P. Watts, Newbury, & Sulentic, 1996) investigated the effects of repeated ascents of a sport climbing route in a sample of 11 expert climbers. The intensity was maximal and pre-post measurements of blood lactate, handgrip strength and endurance were taken. The authors reported correlations between variation in performance in handgrip tests, climbing time and number of laps completed. Variation in blood lactate was only correlated to handgrip strength decreases. Both handgrip strength and endurance remained depressed after 20 minutes of complete rest. Related to this, (Schoffl, Mockel, Kostermeyer, Roloff, & Kupper, 2006) reported lactate accumulations after maximal climbing specific tests on a climbing treadmill. Results were lower ( $5.0 \pm 1.3$  mmol.l<sup>-1</sup>) than in common endurance sports such as running or cycling mainly due to utilization of smaller muscle groups in climbing activities. The researchers also reported that climbers could reach maximal heart rate values during maximal tests performed on a treadmill and the heart rate drop during the rest could explain the recovery ability specific of the sport.

As to no significant results, (Grant et al., 2003) compared climbing specific finger strength endurance in amateur climbers, rowers and aerobically leg trained athletes. No significant differences were seen between the study groups although self-selection was reported as a confounding factor. (Cutts & Bollen, 1993) investigated the differences in pinch and whole hand grip strength between competition climbers and matched controls. Although the climbers' group reported

significantly higher levels of strength, its levels didn't correlate with climbing ability. Only pinch grip strength increased with climbing experience.

Several researchers (MacLeod et al., 2007) studied the factors that could explain the endurance performance in intermediate rock climbers when compared with a matched control group. The researchers found that muscle re-oxygenation during rest periods could predict climbing endurance. These results have been reported in further research (Simon Fryer et al., 2015) that supports the role of re-oxygenation even in the absence of an increased blood flow. Re-oxygenation can be improved through hand shaking during rest periods, with results varying between boulderers and sport climbers due to the length of the rest and activity times (Balas et al., 2016).

A better re-oxygenation during rest periods can result in a decrease in strength loss in the finger flexor muscles, which are crucial for the climbing ability (Philippe, Wegst, Muller, Raschner, & Burtscher, 2012) and have been shown to be stronger in climbers than in non-climbing populations (Vigouroux, Goislard de Monsabert, & Berton, 2015). Fatigue previous to the effort in finger flexors have been shown to impact climbing performance more than fatigue in other muscle groups such as lumbar flexors or shoulder adductors (Deyhle et al., 2015). Finally, forearm flexor oxidative capacity has been reported as one of the best predictors of climbing performance in elite athletes (Simon Fryer et al., 2016) while no differences in oxidative capacities exist between boulderers and sport climbers (S Fryer et al., 2017).

Finally, elite climbers have been reported to have better load balance application than beginners (Donath, Roesner, Schoffl, & Gabriel, 2013).

### ***Climbing and psychological factors***

The first study reviewed (Asci, Demirhan, & Dinc, 2007) examined diverse psychological determinants such as sensation seeking, ability self-perception and intrinsic/extrinsic motivation in a group of Turkish climbers divided by their experience and performance level. The study concluded that climbers had internal motivation, positive self-perception of their abilities and high sensation seeking independently of their experience and current level. Another study (Hardy & Hutchinson, 2007) found a positive correlation between levels of anxiety and performance in climbers. Even in the same subjects, a higher effort was also correlated to increased levels of anxiety. Contrarily, anxiety in beginners has been linked to poorer performances (Nieuwenhuys, Pijpers, Oudejans, & Bakker, 2008). Route previsualization influences climbing style and has been shown to benefit mainly the expert climbers (Sanchez, Lambert, Jones, & Llewellyn, 2012). Higher level climbers report also better stress control, which results in similar performances between top-rope and lead climbing, an equality that is not found in beginners or amateur populations (S Fryer, Dickson, Draper, Blackwell, & Hillier, 2013).

## **DISCUSSION**

This is the first systematic review about performance factors in sport climbing and bouldering. The main findings of this review were:

1. Climbing has moderate cardiorespiratory demands, with reached VO<sub>2</sub>max levels of 30-40 ml/kg/min, blood lactate levels of (5 to 10 mmol/l) and submaximal heart rate (70 to 85% of the maximal heart rate). Thus, even elite climbers don't show high cardiorespiratory capacity.
2. Climbers need significant upper body strength in general and in forearm and finger flexors in particular. Re-oxygenation capacity even in the absence of blood flow in these muscles seems to be the performance limiting factor in repeated intervals of exercise.
3. Evidence shows that anthropometric factors, despite what was previously believed have only a small role in climbing performance. Amongst them, weight seems the most relevant factor due to its role in the relative strength capacity, as it has been demonstrated in hand grip tests.
4. Only a few studies have investigated psychological performance factors in climbing. The most important findings are that route pre-visualization may favor experienced climbers who also show better stress control than beginners.

Figure 2 shows the main performance factors evaluated during the studies that compose this systematic review.



### **Practical applications**

The results obtained in this systematic review could be used in order to design specific training programs based on scientific evidence especially given the future nature of climbing as an Olympic sport. On the other hand, comparative studies between elite and beginners have shown that the non-trainable factors could be used as a potential tool for selection of promising athletes. This systematic review was not designed to analyze the research around nutritional status and climbing performance although the authors acknowledge the importance of dietary manipulation and supplementing (Cabañes, Salinero, Coso, José, & Martín, 2013) in order to change body composition and boost specific energy source utilization (Potter & Fuller, 2015).

### **Future research**

In the following years, studies should focus on correctly assessing training loads in climbing. Until now, this has been done using subjective effort scales and methods that have derived from maximal repetition scales used in strength training (Michailov, 2013). The slow but steady appearance of power measurement tools in sports such as cycling or running may represent future directions for workload assessment in sport climbing and bouldering (Smith & Bird, 2001).

Correct differentiation between beginner, intermediate and elite groups should also be emphasized in a sport that is in constant evolution (N Draper et al., 2011). As new tools have emerged in order to facilitate transfer from climbing grades to statistical analyses and grouping divisions (Nick Draper et al., 2015), future research should correctly assess real performance levels of climbers included in comparative studies and therefore properly establish differences between true elite and advanced or beginner climbers.

### **Limitations**

This systematic review originated from the need of gathering all the scientific research around climbing and performance factors and reviewing all the most important information in order to provide a science-based document for coaches, athletes and researchers around this subject. Thus, a great variability of studies with heterogenous data were included.

The current systematic review excluded non-English and non-Spanish publications; therefore, a possible language bias could appear. The main limitation of this systematic review is the incomplete data provided by the range of heterogeneous studies analyzed during the review: sample sizes were small, performance assessment differed between studies and length of interventions varied substantially, besides the fact that the objectives varied considerably between studies.

## **CONCLUSIONS**

---

Climbing is an intermittent sport that is characterized by a constant fight against gravity. Several trainable performance factors such as upper body strength, specific forearm and finger flexor strength, reoxygenation capacity in the local muscles and power to weight ratio can be seen consistently among elite climbers. Psychological factors such as previsualization capacity or stress control have been described although the research around these subjects is scarce. Anthropometric factors have a smaller role than usually described in common literature. Cardiorespiratory fitness among elite climbers is moderate and similar to the values obtained in other intermittent sports.

## **REFERENCIAS**

---

1. Aras, D., & Akalan, C. (2016). Sport climbing as a means to improve health-related physical fitness parameters. *The Journal of Sports Medicine and Physical Fitness*, 56(11), 1304-1310.
2. Asci, F. H., Demirhan, G., & Dinc, S. C. (2007). Psychological profile of Turkish rock climbers: an examination of climbing experience and route difficulty. *Perceptual and Motor Skills*, 104(3 Pt 1), 892-900. <https://doi.org/10.2466/pms.104.3.892-900>
3. Balas, J., Michailov, M., Giles, D., Kodejska, J., Panackova, M., & Fryer, S. (2016). Active recovery of the finger flexors enhances intermittent handgrip performance in rock climbers. *European Journal of Sport Science*, 16(7), 764-772. <https://doi.org/10.1080/17461391.2015.1119198>
4. Balas, J., Panackova, M., Strejcova, B., Martin, A. J., Cochrane, D. J., Kalab, M., ... Draper, N. (2014). The relationship between climbing ability and physiological responses to rock climbing. *TheScientificWorldJournal*, 2014, 678387. <https://doi.org/10.1155/2014/678387>
5. Bertuzzi, R. C. de M., Franchini, E., Kokubun, E., & Kiss, M. A. P. D. M. (2007). Energy system contributions in indoor rock climbing. *European Journal of Applied Physiology*, 101(3), 293-300. <https://doi.org/10.1007/s00421-007-0501-0>
6. Bertuzzi, R., Franchini, E., Tricoli, V., Lima-Silva, A. E., Pires, F. D. O., Okuno, N. M., & Kiss, M. A. P. D. M. (2012). Fit-climbing

test: a field test for indoor rock climbing. *Journal of Strength and Conditioning Research*, 26(6), 1558-1563. <https://doi.org/10.1519/JSC.0b013e318231ab37>

7. Billat, V., Palleja, P., Charlaix, T., Rizzardo, P., & Janel, N. (1995). Energy specificity of rock climbing and aerobic capacity in competitive sport rock climbers. *The Journal of Sports Medicine and Physical Fitness*, 35(1), 20-24.
8. Cabañes, A., Salinero, J. J., Coso, J. Del, José, J., & Martín, S. (2013). La ingestión de una bebida energética con cafeína mejora la fuerza-resistencia y el rendimiento en escalada deportiva. *Arch Med Deporte*, 30(4), 215-220.
9. Cutts, A., & Bollen, S. R. (1993). Grip strength and endurance in rock climbers. *Proceedings of the Institution of Mechanical Engineers. Part H, Journal of Engineering in Medicine*, 207(2), 87-92. [https:// doi.org/Strength, endurance and anthropometric factors in climbing](https://doi.org/Strength, endurance and anthropometric factors in climbing)
10. Deyhle, M. R., Hsu, H.-S., Fairfield, T. J., Cadez-Schmidt, T. L., Gurney, B. A., & Mermier, C. M. (2015). Relative Importance of Four Muscle Groups for Indoor Rock Climbing Performance. *Journal of Strength and Conditioning Research*, 29(7), 2006-2014. [https:// doi.org/10.1519/JSC.0000000000000823](https://doi.org/10.1519/JSC.0000000000000823)
11. Donath, L., Roesner, K., Schoffl, V., & Gabriel, H. H. W. (2013). Work-relief ratios and imbalances of load application in sport climbing: another link to overuse-induced injuries? *Scandinavian Journal of Medicine & Science in Sports*, 23(4), 406-414. [https:// doi.org/10.1111/j.1600-0838.2011.01399.x](https://doi.org/10.1111/j.1600-0838.2011.01399.x)
12. Draper, N, Dickson, T., Blackwell, G., Priestley, S., Fryer, S., Marshall, H., ... Ellis, G. (2011). Sport-specific power assessment for rock climbing. *The Journal of Sports Medicine and Physical Fitness*, 51(3), 417-425.
13. Draper, Nick, Giles, D., Schöffl, V., Konstantin Fuss, F., Watts, P., Wolf, P., ... Abreu, E. (2015). Comparative grading scales, statistical analyses, climber descriptors and ability grouping: International Rock Climbing Research Association position statement. *Sports Technology*, 8(3-4), 88-94. [https:// doi.org/10.1080/19346182.2015.1107081](https://doi.org/10.1080/19346182.2015.1107081)
14. Egocheaga, J., Montoliu, M., González, V., Rodríguez, B., Del Valle, M., & Palenciano, L. (2001). Metabolismo energético en la escalada deportiva sobre roca y rocódromo versus escalada sobre cascadas de hielo. *Archivos de Medicina Del Deporte*, XVIII(81), 33-40.
15. Fryer, S, Dickson, T., Draper, N., Blackwell, G., & Hillier, S. (2013). A psychophysiological comparison of on-sight lead and top rope ascents in advanced rock climbers. *Scandinavian Journal of Medicine & Science in Sports*, 23(5), 645-650. [https:// doi.org/10.1111/j.1600-0838.2011.01432.x](https://doi.org/10.1111/j.1600-0838.2011.01432.x)
16. Fryer, S, Stone, K. J., Sveen, J., Dickson, T., Espana-Romero, V., Giles, D., ... Draper, N. (2017). Differences in forearm strength, endurance, and hemodynamic kinetics between male boulderers and lead rock climbers. *European Journal of Sport Science*, 17(9), 1177-1183. [https:// doi.org/10.1080/17461391.2017.1353135](https://doi.org/10.1080/17461391.2017.1353135)
17. Fryer, Simon, Stoner, L., Scarrott, C., Lucero, A., Witter, T., Love, R., ... Draper, N. (2015). Forearm oxygenation and blood flow kinetics during a sustained contraction in multiple ability groups of rock climbers. *Journal of Sports Sciences*, 33(5), 518-526. [https:// doi.org/10.1080/02640414.2014.949828](https://doi.org/10.1080/02640414.2014.949828)
18. Fryer, Simon, Stoner, L., Stone, K., Giles, D., Sveen, J., Garrido, I., & Espana-Romero, V. (2016). Forearm muscle oxidative capacity index predicts sport rock-climbing performance. *European Journal of Applied Physiology*, 116(8), 1479-1484. [https:// doi.org/10.1007/s00421-016-3403-1](https://doi.org/10.1007/s00421-016-3403-1)
19. Giles, L. V, Rhodes, E. C., & Taunton, J. E. (2006). The physiology of rock climbing. *Sports Medicine (Auckland, N.Z.)*, 36(6), 529-545. [https:// doi.org/10.2165/00007256-200636060-00006](https://doi.org/10.2165/00007256-200636060-00006)
20. Grant, S., Hasler, T., Davies, C., Aitchison, T. C., Wilson, J., & Whittaker, A. (2001). A comparison of the anthropometric, strength, endurance and flexibility characteristics of female elite and recreational climbers and non-climbers. *Journal of Sports Sciences*, 19(7), 499-505. [https:// doi.org/10.1080/026404101750238953](https://doi.org/10.1080/026404101750238953)
21. Grant, S., Hynes, V., Whittaker, A., & Aitchison, T. (1996). Anthropometric, strength, endurance and flexibility characteristics of elite and recreational climbers. *Journal of Sports Sciences*, 14(4), 301-309. [https:// doi.org/10.1080/02640419608727715](https://doi.org/10.1080/02640419608727715)
22. Grant, S., Shields, C., Fitzpatrick, V., Loh, W. M., Whitaker, A., Watt, I., & Kay, J. W. (2003). Climbing-specific finger endurance: a comparative study of intermediate rock climbers, rowers and aerobically trained individuals. *Journal of Sports Sciences*, 21(8), 621-630. [https:// doi.org/10.1080/0264041031000101953](https://doi.org/10.1080/0264041031000101953)
23. Hardy, L., & Hutchinson, A. (2007). Effects of performance anxiety on effort and performance in rock climbing: a test of processing efficiency theory. *Anxiety, Stress, and Coping*, 20(2), 147-161. [https:// doi.org/10.1080/10615800701217035](https://doi.org/10.1080/10615800701217035)
24. La Torre, A., Crespi, D., Serpiello, F. R., & Merati, G. (2009). Heart rate and blood lactate evaluation in bouldering elite athletes. *The Journal of Sports Medicine and Physical Fitness*, 49(1), 19-24.
25. Laffaye, G., Levernier, G., & Collin, J.-M. (2016). Determinant factors in climbing ability: Influence of strength, anthropometry, and neuromuscular fatigue. *Scandinavian Journal of Medicine & Science in Sports*, 26(10), 1151-1159. [https:// doi.org/10.1111/sms.12558](https://doi.org/10.1111/sms.12558)
26. Limonta, E., Brighenti, A., Rampichini, S., Cè, E., Schena, F., & Esposito, F. (2018). Cardiovascular and metabolic responses during indoor climbing and laboratory cycling exercise in advanced and elite climbers. *European Journal of Applied Physiology*, 118(2), 371-379. [https:// doi.org/10.1007/s00421-017-3779-6](https://doi.org/10.1007/s00421-017-3779-6)
27. Lutter, C., El-Sheikh, Y., Schöffl, I., & Schöffl, V. (2017). Sport climbing: medical considerations for this new Olympic discipline. *British Journal of Sports Medicine*, 51(1), 2-3. [https:// doi.org/10.1136/bjsports-2016-096871](https://doi.org/10.1136/bjsports-2016-096871)
28. MacLeod, D., Sutherland, D. L., Buntin, L., Whitaker, A., Aitchison, T., Watt, I., ... Grant, S. (2007). Physiological determinants of climbing-specific finger endurance and sport rock climbing performance. *Journal of Sports Sciences*, 25(12), 1433-1443. [https:// doi.org/10.1080/02640410600944550](https://doi.org/10.1080/02640410600944550)
29. Mermier, C. M., Janot, J. M., Parker, D. L., & Swan, J. G. (2000). Physiological and anthropometric determinants of sport climbing performance. *British Journal of Sports Medicine*, 34(5), 359-365; discussion 366.
30. Mermier, C. M., Robergs, R. A., McMinn, S. M., & Heyward, V. H. (1997). Energy expenditure and physiological responses during indoor rock climbing. *British Journal of Sports Medicine*, 31(3), 224-228.
31. Michailov, M. L. (2013). Workload characteristic, performance limiting factors and methods for strength and endurance training in rock climbing. *Med Sport*, (18), 97-106. [https:// doi.org/10.5604/17342260.1120661](https://doi.org/10.5604/17342260.1120661)

32. Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G., & PRISMA Group. (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Medicine*, 6(7), e1000097. [https:// doi.org/10.1371/journal.pmed.1000097](https://doi.org/10.1371/journal.pmed.1000097)
33. Nieuwenhuys, A., Pijpers, J. R., Oudejans, R. R. D., & Bakker, F. C. (2008). The influence of anxiety on visual attention in climbing. *Journal of Sport & Exercise Psychology*, 30(2), 171-185.
34. Ozimek, M., Rokowski, R., Draga, P., Ljakh, V., Ambrozy, T., Krawczyk, M., ... Mucha, D. (2017). The role of physique, strength and endurance in the achievements of elite climbers. *PloS One*, 12(8), e0182026. [https:// doi.org/10.1371/journal.pone.0182026](https://doi.org/10.1371/journal.pone.0182026)
35. Philippe, M., Wegst, D., Muller, T., Raschner, C., & Burtcher, M. (2012). Climbing-specific finger flexor performance and forearm muscle oxygenation in elite male and female sport climbers. *European Journal of Applied Physiology*, 112(8), 2839-2847. [https:// doi.org/10.1007/s00421-011-2260-1](https://doi.org/10.1007/s00421-011-2260-1)
36. Potter, J., & Fuller, B. (2015). The effectiveness of chocolate milk as a post-climbing recovery aid. *The Journal of Sports Medicine and Physical Fitness*, 55(12), 1438-1444.
37. Rodio, A., Fattorini, L., Rosponi, A., Quattrini, F. M., & Marchetti, M. (2008). Physiological adaptation in noncompetitive rock climbers: good for aerobic fitness? *Journal of Strength and Conditioning Research*, 22(2), 359-364. [https:// doi.org/10.1519/JSC.0b013e3181635cd0](https://doi.org/10.1519/JSC.0b013e3181635cd0)
38. Sanchez, X., Lambert, P., Jones, G., & Llewellyn, D. J. (2012). Efficacy of pre-ascent climbing route visual inspection in indoor sport climbing. *Scandinavian Journal of Medicine & Science in Sports*, 22(1), 67-72. [https:// doi.org/10.1111/j.1600-0838.2010.01151.x](https://doi.org/10.1111/j.1600-0838.2010.01151.x)
39. Schoffl, V. R., Mockel, F., Kostermeyer, G., Roloff, I., & Kupper, T. (2006). Development of a performance diagnosis of the anaerobic strength endurance of the forearm flexor muscles in sport climbing. *International Journal of Sports Medicine*, 27(3), 205-211. [https:// doi.org/10.1055/s-2005-837622](https://doi.org/10.1055/s-2005-837622)
40. Schreiber, T., Allenspach, P., Seifert, B., & Schweizer, A. (2015). Connective tissue adaptations in the fingers of performance sport climbers. *European Journal of Sport Science*, 15(8), 696-702. [https:// doi.org/10.1080/17461391.2015.1048747](https://doi.org/10.1080/17461391.2015.1048747)
41. Smith, M. F., & Bird, S. R. (2001). Reliability of Mean Power Recorded During Indoor and Outdoor Self-Paced 40 km Cycling Time-Trials Exercise Supervision in the Workplace View project Assessing the effects of regional diets on Cardiovascular Disease Risk reduction View project. *Article in International Journal of Sports Medicine*. [https:// doi.org/10.1055/s-2001-13813](https://doi.org/10.1055/s-2001-13813)
42. Vigouroux, L., Goislard de Monsabert, B., & Berton, E. (2015). Estimation of hand and wrist muscle capacities in rock climbers. *European Journal of Applied Physiology*, 115(5), 947-957. [https:// doi.org/10.1007/s00421-014-3076-6](https://doi.org/10.1007/s00421-014-3076-6)
43. Wall, C. B., Starek, J. E., Fleck, S. J., & Byrnes, W. C. (2004). Prediction of indoor climbing performance in women rock climbers. *Journal of Strength and Conditioning Research*, 18(1), 77-83.
44. Watts, P., Newbury, V., & Sulentic, J. (1996). Acute changes in handgrip strength, endurance, and blood lactate with sustained sport rock climbing. *The Journal of Sports Medicine and Physical Fitness*, 36(4), 255-260.
45. Watts, P B, Daggett, M., Gallagher, P., & Wilkins, B. (2000). Metabolic response during sport rock climbing and the effects of active versus passive recovery. *International Journal of Sports Medicine*, 21(3), 185-190. [https:// doi.org/10.1055/s-2000-302](https://doi.org/10.1055/s-2000-302)
46. Watts, Phillip B. (2004). Physiology of difficult rock climbing. *European Journal of Applied Physiology*, 91(4), 361-372. [https:// doi.org/10.1007/s00421-003-1036-7](https://doi.org/10.1007/s00421-003-1036-7)
47. White, D. J., & Olsen, P. D. (2010). A time motion analysis of bouldering style competitive rock climbing. *Journal of Strength and Conditioning Research*, 24(5), 1356-1360. [https:// doi.org/10.1519/JSC.0b013e3181cf75bd](https://doi.org/10.1519/JSC.0b013e3181cf75bd)